



International Journal of Pharmaceutical Research and Development (IJPRD)

Platform for Pharmaceutical Researches & Ideas

www.ijprd.com

SPECTROPHOTOMETRIC METHOD OF ESTIMATION OF SIMVASTATIN BY SULFO-PHOSPHO-VANILLIN REACTION

Singh Smita^{1*}

¹Kalka group of institution research and advanced studies, Meerut (U. P.)

ABSTRACT

The importance of the quinolone as a nucleus with medicinal properties is well established, since many decades. The nucleus possesses pharmacological activities such as antibacterial antimicrobial, anticonvulsant activity and antiHIV. This nucleus possess anticonvulsant activity that why this property is use for the treatment that are related to mental disorder, like seizures. Quinolone and its derivatives possessing triazolone and triazolo nucleus has attracted great attention in recent years due to wide variety of biological activity particularly anticonvulsant activity keeping in view the continuing interest in quinolone and its derivatives, the present study is aimed at synthesizing safer and effective quinolone derivatives by using microwave.

Key words: quinolone, triazolo, synthesis, antibacterial activity.

Correspondence to Author



SINGH SMITA

Kalka group of institution research and advanced studies, Meerut (U. P.)

Email: smita0082@gmail.com

INTRODUCTION

The rapid rise in microbial resistance to the traditional antibiotics has necessitated a continuing search for new classes of compounds with novel modes of antibacterial activity. The 4-quinolone antibacterial has a number of advantages over other classes of antibacterial agents. They have wide spectrum of activity, well absorbed orally, have relatively long serum half lives and minimal toxicity. The currently used quinolone derivatives are known to have several drug interactions and adverse side effects.(ramiaha et al.,)

The chemicals and reagents used in the present project were of AR and LR grade, procured from Sigma Aldrich, Aventis Pharma limited Goa, Qualigens and Ranchem. Melting points of the synthesized compounds were determined by open capillary method. The IR spectra of the synthesized compounds were recorded on a Fourier Transform IR spectrometer (model Shimadzu 8400S) at Shobhit University, school of pharmaceutical sciences Meerut. ¹H NMR spectra were recorded on amx-400 NMR spectrometer at Punjab University using D₂O and chemical shifts (δ) are reported in parts per million downfield from internal reference Tetramethylsilane (TMS). Mass

EXPERIMENTAL WORK

MATERIAL (S) AND METHOD (S)

spectra were obtained from Punjab University and recorded on Shimadzu LC-MS model 2010A.

General procedure for the synthesis of 4-quinolone-3-carboxylic acid:-

General procedure

A mixture of the appropriate aniline (1 mmol) and diethyl ethoxy methylene manolate (200µml, 1mmol) and polyphosphoric acid was kept in microwave at moderate temperature for 1.5 minute. The crystal solid was obtained after cooling to room temperature.

A solid of the appropriate ester (1mmol) in 10% aq.NaOH (10ml) was reflux for 2 hr. After cooling at room temperature, the reaction mixture was acidified using conc. HCl. The resulting precipitate was filtered and washed with water to give the corresponding 4-quinolone-3-carboxylic acid.

Experimental procedure

Synthesis of 4-quinolone-3-carboxylic acid derivatives (1)

A mixture of the 2-amino 5-Bromo benzoic acid (1 mmol) and diethyl ethoxy methylene manolate (200µml, 1mmol) and polyphosphoric acid was in microwave at moderate temperature for 1.5 minute. The crystal solid was obtained after cooling to room temperature. **(Step 1)**

S.NO.	R ₁	R ₂
1	-Br	-COOH
2	-CH ₃	-H
3	-OH	-H

A solid of 6-bromo 4-oxo-1,4 dihydro quinolone 3,8 dicarboxylic acid 3 ethyl ester (1mmol) in 10% aq.NaOH (10ml) was reflux for 2 hr. After cooling at room temperature, the reaction mixture was acidified using conc. HCl. The resulting precipitate was filtered and washed with water to give the corresponding 4-quinolone-3-carboxylic acid. The yields of the products were between 62-70 %. **(Step 2)**

Synthesis of 4-quinolone-3-carboxylic acid derivatives (2)

A mixture of the 4-amino-phenol (1 mmol) and diethyl ethoxy methylene manolate (200µml, Available online on www.ijprd.com

1mmol) and polyphosphoric acid was in microwave at moderate temperature for 1.5 minute. The crystal solid was obtained after cooling to room temperature add equal amount of distilled water and recrystallized by DMF. **(Step 1)**

A solid of the 6-hydroxy 4-oxo-1, 4 dihydro quinolone 3-carboxylic acid ethyl ester (1mmol) in 10% aq.NaOH (10ml) was reflux for 2 hr. After cooling at room temperature, the reaction mixture was acidified using conc. HCl. The resulting precipitate was filtered and washed with water to give the corresponding 4-quinolone-3-carboxylic acid. **(Step 2)**

Synthesis of 4-quinolone-3-carboxylic acid derivatives (3)

A mixture of the p-toluidine (1 mmol) and diethyl ethoxy methylene manolate (200µml, 1mmol) and polyphosphoric acid was in microwave at moderate temperature for 1.5 minute. The crystal solid was obtained after cooling to room temperature add equal amount of distilled water and recrystallized by DMF. **(Step 1)**

A solid of the 6-methyl 4-oxo-1, 4 dihydro quinolone 3 carboxylic acid ethyl ester (1mmol) in 10% aq.NaOH (10ml) was reflux for 2 hr. After cooling at room temperature, the reaction mixture was acidified using conc. HCl. The resulting precipitate was filtered and washed with water to give the corresponding 4-quinolone-3-carboxylic acid. **(Step 2)**

S.NO.	R ₁	R ₂
1	-Br	-COOH
2	-CH ₃	H-
3	-OH	H-

IR Spectral analysis of synthesized compounds

FTIR Spectra of compounds were recorded using potassium bromide pellets (Perkin Elmer spectrum, RXI FT-IR System). Sample and potassium bromide (IR grade) was mixed in 1:100 ratios and scanning was done between 4000-400 cm⁻¹. Interpretation analyses of synthesized derivatives are summarized in tables.

Antibacterial activity:

Antibacterial activity of the synthesized compounds was determined by the serial dilution method against the gram-positive organisms *Bacillus subtilis* and gram-negative organisms *Escherichia coli*, species at different concentration. The bacteria were subcultured on Nutrient Agar

medium. The petri dishes were incubated at 37°C for 24h. Standard antibacterial drugs were also screened under similar conditions for comparison. 0.1ml of Ciprofloxacin (300µg/ml) (Std.) was used as standard for other microorganisms. The results are presented in Table(a).

Table (a) Physical parameters of synthesized derivatives-

S.NO.	COMPOUND	% yield	Melting range (°C)	R _f value	R1	R2
1.	1	60-70	138-140	0.71	-Br	-COOH
2.	2	50-60	163-165	0.32	-CH ₃	-H
3.	3	50-60	185-187	0.34	-OH	-H

RESULTS AND DISCUSSION

Quinolin-4-one-3-carboxylic acid derivatives were synthesized and screened for their efficacy as antibacterial agents against various pathogens in-vitro by Agar dilution method. The stock solution (100mg/ml) of Quinolin-4-one-3-carboxylic acid derivatives was prepared in dimethyl sulphoxide and working solution of concentrations 1, 2, 4, 8, 16 and 32 µg/ml in distilled water was prepared. Same dilutions were prepared for the standard drug.

Inhibition of Escherichia coli

All the synthesized compounds (1-3) showed inhibition of E.coli at concentration range from 2µg/ml to 16µg/ml. The compound **2** showed higher activity (less MIC) when compared with standard drug at concentration of 2µg/ml. The compound **3** showed equipotent activity with standard drug at 4µg/ml.

Table (b) Antibacterial activity of compounds (1-3)

S.NO.	(MIC-µg/ml)		
	Compound Code	Escherichia coli	Bacillus subtilis
1	1	2	8
2	2	4	16
3	3	8	16
Standard drug	Ciprofloxacin	4	8

Standard Drug (Ciprofloxacin (100µg/mL))

ACKNOWLEDGEMENT

The authors thank **Miss. Sarika Sahu**, for her help during microbiological work. Their thanks are also due to **Mr. Parag** for providing the cultures of micro-organisms and chemicals, Punjab University, for the ¹HNMR reports and also for the Mass spectral reports. They also wish to express their thanks to the management, director (**Prof. Ranjit Singh**) and my guide (**Mr. Anand Gaurav**) of Shobhit University for the facilities provided.

REFERENCES

1. Roy, A.; Sardar, S. M.; Salve, B.U. Synthesis some novel fluoroquinolone derivatives and evaluated them for antibacterial activity. *International Journal of Chem Tech Research* **2009**, *1* (1), 34-39.
2. Yusun, X.; Zhang, L.; Quan, Z. S.; Piao, H. R.; Wei, C. X. Design, synthesis 8 alkoxy-5, 6 dihydro-[1, 2, 4] triazino [4, 3-a] Quinoline-1-ones and their anticonvulsant activity. *European Journal of Medicinal Chemistry* **2009**, *44*, 1265-1271.
3. Tzeng, C. C.; Chen, Y. L.; Chen, Y. W.; Lo, W. F.; Kao, C. L. Synthesis aryl sulfonyl and Aryl carbonyl derivatives of ofloxacin and evaluated them for antimycobacterial activity. *Journal of Chinese Chemical Society* **2009**, *56*, 374-380.
4. Li, X.; Huang, X.; Ningwu, C. Synthesis and evaluation of a series of a novel C7 non-basic substituted fluoroquinolones as antibacterial agents *Bioorganic & Medicinal Chemistry Letters* **2009**, *19*, 4130-4138.
5. Shafiee, A.; Foroumadi, A. Synthesis some new N-piperazinyl quinolone derivatives and evaluated them for antibacterial activity. *J. Iran. Chem. Soc.* **2009**, *6*(2), 325-330.
6. Illig, C.R.; Wall, M. J.; Chen, J.; Meegalla, S.; Wilson, K.J. Synthesis some novel 3, 4, 6-substituted 2-quinolones and evaluated them as FMS kinase inhibitors. *Bioorganic & Medicinal Chemistry Letters* **2008**, *18*, 2097-2102.
7. You, Q. D.; Wang, H. Design, Synthesis and evaluation of antitumor activity of a series of 3-substituted quinolone derivatives. *Chinese Chemical Letters* **2008**, *19*, 1395-1401.
8. Corelli, F.; Pasquini, S.; Mugnaini, C. Synthesis of 4-quinolone-3-carboxylic acid derivatives and evaluated them for anti-HIV activity. *J. Med. Chem* **2008**, *51*, 5125-5130.
9. Shafie., A.; Sorkhi, M.; Synthesis some novel N-[2-(2-naphthyl) ethyl] piperazinyl quinolone derivatives and evaluation for their antibacterial activity. *DARU* **2008**, *16*(3), 189-195.
10. Nagao, Y.; Ikee, Y.; Hashimoto, K.; Nakashima, M. Synthesis new quinolone derivatives utilizing 1-azabicyclo [1.1.0] butane and evaluated them for antibacterial activity. *Bioorganic & Medicinal Chemistry Letters* **2007**, *17*, 942-945.
11. Souza, M. L. D.; Ferreira, V. F.; Lucero, B. A.; Gomes, C. R. B. Synthesis several 1-[(2-hydroxy-ethoxy) methyl]-3-carbethoxy-4(1H) quinolones and 1-[(2-hydroxy-ethoxy) methyl]-4(1H) quinolone-3-carboxylic acids as quinolonic acyclovir analogues and evaluated anti-HSV-1 activity against herpes simplex virus type 1 (HSV-1). *Bioorganic & Medicinal Chemistry Letters* **2006**, *16*, 1010-1013.
12. Yusun, X.; Zhe Jin, Y.; Nahli, F.; Yunchai, K.; Quan, Z. S.; Y. Shanquan, Z. Synthesis of 8 alkoxy-4, 5, dihydro-[1, 2, 4] triazino [4, 3-a] Quinoline-1-ones and evaluation for their anticonvulsant activity. *Arch Pharm Res* **2006**, *29*, 1080-1086.
13. He, Y.; Ellis, D.; Kuhlen, K.L.; Anaclerio, B.; Wu, B.; Yin, H.; Bursulaya, B. Synthesis and design novel quinolone derivatives and evaluation for their as HIV-1 non-nucleoside reverse transcriptase inhibitors. *Bioorganic & Medicinal Chemistry Letters* **2006**, *16*, 4246-4251.
14. Asahina, Y.; Iwase, K.; Ishiaki, T.; Hosaka, M.; linuma, F. Synthesis of 1-(2-Fluorovinyl)-7-substituted-4-quinolone-3-carboxylic acid derivatives and evaluation for their antibacterial activity. *J. Med. Chem* **2005**, *48*, 3194-3202.
15. Foroumadi, A.; Shafiee, A.; Emami, S.; Rajaei, M. Synthesis of some novel N-(5-benzylthio-1,

- 3, 4-thiadiazol-2-yl) and N-(5-benylsulfonyl-1, 3, 4-thiadiazol-2-yl) piperazinyl quinolone derivatives, evaluation for antibacterial activity. *Bioorganic & Medicinal Chemistry Letters* **2005**, *15*, 4488-4492.
16. Asahina, Y.; Hosaka, M.; Iwase, K.; Ishizaki, T. Synthesis of some novel 4-quinolone-3-carboxylic acid derivatives having a trifluoromethyl group as a novel N-1 substituent, and evaluated them for antibacterial activity. *J. Med. Chem* **2005**, *48*, 3443-3450.
17. Park, M. S. Inlee, J.; Sookpark, M. Synthesis some novel 2, 3-dihydro-2-phenyl-4-quinolone derivatives. *Bull. Korean Chem. Soc* **2004**, *25*(8), 1269-1272.
18. Guo, H.Y.; Liu, J.Y.; Lu, Y.; Wang, Y. C. Synthesis a series of 5-amino-8-methoxyquinolones derivatives and evaluation them for their antibacterial activity. *Chinese Chemical Letters* **2004**, *15*(12), 1397-1402.
19. Sarkozy, G. Quinolones: a class of antimicrobial agents. *Vet. Med-Czech* **2001**, *46*, 257-274.
20. Jaradat, N. A.; Taran, S. G.; Likhanova, N.V.; Gorokhova, O. V.; Ukrainets, I.V. Synthesis, structurally analysis and anticonvulsant activity of optically active 3-quinoline carboxylic acids derivatives. *Chemistry of Heterocyclic Compounds* **2000**, *36*(1), 49-53.
21. Koyama, J.; Toyokuni, I.; Tagahara, K. Synthesis 2-Arylquinoline and 2-Aryl-4-quinolone Alkaloids Diels–Alder Reaction of 1, 2, 3-Benzotriazine with Enamines.
22. Kuo, C. S.; Jiauhuang, Li.; Chiehhsieh, M.; Mingteng, C. Synthesis of phenyl quinolones evaluation for antiplatelet activity. *Bioorganic & Medicinal Chemistry Letters* **1998**, *6*, 1657-1661.
23. Kidwai, M.; Misra, P.; Kumar, R.; Saxena, R.K. Synthesis by using microwave assisted method and antibacterial activity of new quinolone derivatives. *Monatshefte fur Chemie* **1998**, *129*, 961-965.
24. Tillotson, G. S. Quinolones: structure-activity relationships and future predictions. *J. Med. Microbiol* (**1996**), *4*, 320-325.
25. Batori, S.; Timari, G.; Koczka, I.; Hermecz, I. Synthesis and evaluation of N-(1-aziridino)-6-fluoro-quinolone-3-carboxylic acids. *Bioorganic & Medicinal Chemistry Letters* **1996**, *6*(13), 1507-1513.
26. Stadlbauer, W.; Steinschifter, W. Synthesis of 4-Azido-2(1H)-quinolones by thermolytic degradation of oxazolo [4, 5-c] quinolones. *J. Heterocyclic. Chem* **1994**, *31*, 1647-1651.
27. Kappe, T.; Karem, A. S.; Stadlbauer, W. Synthesis of benzo-halogenated 4-hydroxy-2(1H) quinolones. *J. Heterocyclic chem* **1998**, *25*, 5857-862.
28. Willey, J.M.; Sherwood, L.M.; Woolverton, C.J.; 'Prescott, Harley and Wein's Microbiology', 7th edition, McGraw Hill Higher education. 2001; 835-840.
29. Jain, N.K. Pharmaceutical Microbiology ,2nd edition, Vallabh Prakashan,**1999**; 46-48, 295-300
30. European Committee for Antimicrobial Susceptibility Testing (EUCAST) of the European Society of Clinical Microbiology and Infectious Diseases (ESCMID). Determination of minimum inhibitory concentrations (MICs) of antibacterial agents by agar dilution. Eucast definitive document e.def 3.1 June **2000**, 509-515.
31. Venkataramana, C.H.S.; Gupta. Manish; *International Journal of Pharma Research and Development* synthesis of novel pyrazolidinedione substituted 4-quinolone derivatives and their biological activity.
